



Metrosideros polymorpha ('ōhi'a lehua)

Myrtaceae (myrtle family)

'ōhi'a, 'ōhi'a lehua, lehua (Hawai'i)

J. B. Friday and Darrell A. Herbert

IN BRIEF

Distribution Endemic to the six largest Hawaiian islands.

Size Greatly variable, typically reaches 20–24 m (66–79 ft) in height; may be much smaller on lava rock or boggy soils.

Habitat Wide range, 1–2500 m (1–8200 ft) elevation with rainfall of 400–10,000 mm (16–400 in). Largest component of lowland and montane wet and mesic forests, dry forests, subalpine shrublands, and new lava flows.

Vegetation Associated with dozens of other native species.

Soils Very widely adapted, occurring on medium and heavy clay uplands; tolerates rocky soils and organic soils forming on recent lava flows.

Growth rate Slow growing, 0.3–0.6 m (12–24 in) per year in height, and 1–3 mm (0.04–0.12 in) in stem diameter per year.

Main agroforestry uses Windbreak, shade, ornamental.

Main products Pole wood, fuelwood, honey source

Yields High-volume, old-growth stands may yield 70–84 m³/ha (1000–1200 ft³/ac or 5000–6000 bf/ac); stands on poor soils such as lava rock may yield much less.

Intercropping Used in farm cultivation only when naturally present due to very slow establishment and growth.

Invasive potential Poses a serious risk of being invasive if introduced outside of Hawai'i.



Huge old 'ōhi'a tree at Kīpuka Puaulu, Hawai'i Volcanoes National Park. Note the deeply fluted bole and the sparse crown.

INTRODUCTION

The native Hawaiian ‘ōhi‘a (*Metrosideros polymorpha*) is the most abundant tree in the Hawaiian Islands. The name *Metrosideros* is derived from the Greek *metra*, heartwood, and *sideron*, iron, in reference to the hard wood of the genus (Dawson and Stemmermann 1999). Known locally by its Hawaiian name, ‘ōhi‘a lehua, the species is found on all the major islands and in a variety of habitats. Distributed from near sea level to the tree line (2500 m [8200 ft]), the species exhibits tolerances of frost, volcanic vapors, and excesses or deficiencies in moisture. It is found across a rainfall gradient having extremes of less than 400 mm (16 in) to over 10,000 mm (400 in) on Mt. Wai‘ale‘ale, Kaua‘i, one of the wettest places on earth. ‘Ōhi‘a is found growing on a broad range of substrates including those as young as recent lava flows or as old as highly weathered Oxisols. Dominant in cloud forests and most rainforests above 400 m (1300 ft), often with a tree fern understory, the species is also common in seasonally wet forests, where it may be dominant or in mixtures with the native *Acacia koa* or the invasive species *Morella faya* (formerly *Myrica*) (Mueller-Dombois and Fosberg 1998).

‘Ōhi‘a is a slow-growing broad leaf evergreen whose form, foliage, and flower color can vary greatly from site to site. Representatives of the species can reach a stature as great as 30 m (100 ft) on well drained soils with sufficient moisture, while others in boggy sites can be fully mature when only a few cm tall. This phenotypic variation, affected by environment, allows the species to adjust to the extreme range of conditions listed above (St. John 1979, Stemmermann 1983, Aradhya et al. 1991). The resulting expressions of morphological variation lie at the heart of the species name *polymorpha* (meaning “many forms”) and have prompted investigations into the genetic variations. The species is thought to be an example of how one ancestral species can begin to diversify and evolve into many species when isolated in an island environment (incipient speciation). It may also be that ‘ōhi‘a remains a generalist with broad plasticity.

There are many modern and ancient uses for the flowers, leaves, and wood of ‘ōhi‘a, but the forests are perhaps most highly valued for aesthetics, watershed protection, and as habitat for Hawai‘i’s native and endemic biota. The species itself is a critical food source to endemic nectivorous and insectivorous birds (Berger 1981, Smith et al. 1995), and critical specific habitat to endemic *Achatinella* and *Partulina* tree snail species (Hadfield and Mountain 1980, Hadfield and Miller 1989). However, past land-use practices have resulted in the reduction of ‘ōhi‘a’s range, largely through land-use conversion to cattle pasture and agriculture. Because mature stands may experience canopy decline and dieback of age-cohorts (Mueller-Dombois 1983, 1985, 1987),

Hawai‘i’s early foresters experimented with the replacement of some forest stands with introduced species for the purposes of maintaining watershed integrity and creating a potentially renewable timber industry. In recent years there have been efforts to restore some of these lost ranges.

DISTRIBUTION

The natural distribution is limited to the six largest islands in the Hawaiian archipelago: Hawai‘i, Maui, Moloka‘i, Lāna‘i, O‘ahu, and Kaua‘i.

Location of introductions

‘Ōhi‘a is planted as an ornamental plant in New Zealand, where there are several other native species of *Metrosideros* (Adams 1967).

BOTANICAL DESCRIPTION

Preferred scientific name

Metrosideros polymorpha Gaud.

Family

Myrtaceae (myrtle family)

Non-preferred scientific names

Metrosideros collina (not itself a synonym, but several infraspecific taxa described under it are)

M. haleakalensis

M. lutea

Nania glabrifolia

N. lutea

N. polymorpha

Common names

‘ōhi‘a, ‘ōhi‘a lehua, lehua (Hawai‘i). The common names are often written without the Hawaiian diacritical marks, i.e., ohia or ohia lehua.

Growth, form, and size

Growth form ranges from tall trees to small erect or prostrate shrubs. Crowns may be compact or open-branched. Height of crown base may be low when growing in open habitat, but branching begins at mid-stem or higher in closed canopy forest. On moist, deep soils, ‘ōhi‘a grows to 20–24 m (66–79 ft) in height with a diameter at breast height (dbh) of 45–90 cm (18–36 in). Trees to 30 m (100 ft) in height and 216 cm (85 in) dbh have been recorded (Lamb 1981, Elevitch pers. obs.). The bole may be straight,



Top left: Flower buds close to opening. Top right: Ripening seed capsules; open capsules show tiny seeds within. Bottom left: Jack Jeffrey shows the common red flower, so familiar in the remaining native Hawaiian forests. Bottom right: Typical appearance of bark. PHOTOS: C. ELEVITCH

cylindrical and smooth, or twisted and deeply fluted. Forest trees may often have tall stilt roots, as these trees grow from seedlings that germinate on fallen logs or fallen hāpu‘u (*Cibotium* spp., tree fern) stems, which rot away by the time the tree matures. Some trees may sprout bunches of aerial roots. Shrub growth forms predominate on poorly drained, dry or rocky soils, and at high altitude (Skolmen 1974a, Little and Skolmen 1989, Adee and Conrad 1990, Mueller-Dombois and Fosberg 1998, Dawson and Stemmermann 1999).

Flowers

Flowers are usually red but sometimes salmon, pink, yellow, or orange. Clusters of flowers are about 5–8 cm (2–3 in) wide and are on the terminal ends of the branches. The bunches of stamens extend 1–3 cm (0.4–1.2 in) out from the flower and give the blossoms a pompom, brush, or hair-like appearance. The Hawaiian name *lehua* (from *‘ōhi‘a*

lehua) refers to the hair-like appearance of the blossoms (Rock 1974). Flowers occur in inflorescences of 2–5 pairs of cymules; peduncles 7–18 mm (0.28–0.7 in) long; pedicels 2–8 mm (0.08–0.32 in) long; bracts ovate to suborbicular, 5–10 mm (0.2–0.4 in) long; sepals rounded to triangular, 1.5–4 mm (0.06–0.16 in) long; petals obovate to orbicular, 2.5–5 mm (0.1–0.2 in) long; stamens 10–30 mm (0.4–1.2 in) long; style 13–30 mm (0.5–1.2 in) long (Dawson and Stemmermann 1999). Each cluster of flowers lasts several days. Flowers are pollinated by both birds and insects. Yellow flowers have been shown to be self-compatible, while red flowers are partially self-incompatible and require a pollinator for good seed set (Carpenter 1976).



Pink flowers. PHOTO: J. B. FRIDAY



Yellow flowers. PHOTO: J. B. FRIDAY



Salmon flowers. PHOTO: C. ELEVITCH

Leaves

Foliage varies from thick, dark greenish-gray with dense surface fuzziness, primarily on the lower surface of the leaf,

to thin, bright green, and smooth. Leaves are opposite and clustered at the ends of the branches. Leaf shape is highly variable, ranging from round to oval to egg-shaped, 1–8 cm (0.4–3.1 in) long, 1–5.5 cm (0.4–2.2 in) wide. Several pairs of raised veins radiate from the base. Upper surface is smooth, lower surface smooth or woolly or fuzzy. Margins are flat to turned under, apex rounded or sometimes pointed, base wedge-shaped or heart-shaped. Petioles are 1–16 mm (0.04–0.63 in) long, 1–3 mm (0.04–0.1 in) wide, but can be sessile, especially at high elevation (Dawson and Stemmermann 1999).

Fruit

Fruiting capsules are 3–10 mm (0.1–0.4 in) long and wide, smooth or fuzzy (Dawson and Stemmermann 1999).

Seeds

The seed capsules contain many minute seeds. There are approximately 1750 seeds per gram (50,000 seeds/oz).

Similar or look-alike species

There are about 50 species in the genus *Metrosideros* in Southeast Asia and the Pacific (Wagner et al. 1999). *Metrosideros kermadecensis*, from the Kermadec Islands off of New Zealand, has escaped from cultivation on Maui and may be considered a pest species (Evenhuis and Eldredge 2004). The pohutukawa tree of New Zealand, *Metrosideros excelsa* (syn. *M. tomentosa*), is sometimes planted as an ornamental in Hawai'i but is not reported to have escaped from cultivation (Kinsey, undated). Several cultivars of *M. excelsa* are grown. 'Ōhi'a was originally classified as a variety of *Metrosideros collina*, a related tree of Tahiti, Rarotonga, and other islands in the South Pacific, but now is seen as a species endemic to Hawai'i (Little and Skolmen 1989).

How to distinguish from similar species/look-alikes

The flowers of *Metrosideros excelsa* and *M. kermadecensis* are dark crimson in color, as opposed to the more orange-red of *M. polymorpha*, and the young buds and undersides of leaves are covered in a silvery-white felt. *Metrosideros kermadecensis* is a smaller tree with brighter red flowers than *M. excelsa*. A variegated leaf variety of *M. kermadecensis* is also cultivated. *Metrosideros polymorpha* has not been reported to naturalize elsewhere in the Pacific, although it has been introduced to New Zealand as an ornamental plant. The Hawaiian species of *Metrosideros* are characterized by the presence of bud scales and a frequently forking branching pattern.



Metrosideros kermadecensis, an exotic to Hawai‘i not recommended for planting, has darker crimson leaves than ‘ōhi‘a, and a silvery-white felt on the underside of the leaves. PHOTO: FOREST AND KIM STARR, USGS

GENETICS

Eight varieties of the Hawaiian *Metrosideros polymorpha* complex are recognized on the basis of classical taxonomy (Wagner et al. 1999). In addition to the *M. polymorpha* varieties, four distinct species are also recognized: *M. macropus*, *M. rugosa*, *M. tremuloides*, and *M. waialealae*. Genetic distinctions between varieties are not clear, but isozyme analyses suggest variability between populations are a likely case of incipient speciation (Aradhya et al. 1991). Altitudinal, soil-based, and successional ecotypes have been proposed on the basis of morphological variations, especially in leaves (Corn and Hiesey 1973, Stemmermann 1983, Stemmermann and Ihsle 1993, Adee and Conrad 1990, Mueller-Dombois 1994, Kitayama et al. 1997). Intraspecific hybridization has been demonstrated (Corn 1979). Genetic differences correlate with environmental differences, although in some cases differences within phenotypic varieties may be greater than between varieties, and many intermediate phenotypes exist (James et al. 2004).

Variability of species

The species occurs in a range of forms from prostrate shrubs to tall trees. Leaves range greatly in both size and shape and may be smooth to fuzzy or woolly. Flowers may be red, pink, salmon, orange, yellow, and many shades in between.

Known varieties

There are eight known varieties (see table), some of which

<i>dieteri</i>	Middle to high elevations on Kaua‘i. Small trees. Leaves with raised veins.
<i>glaberrima</i>	Middle to high elevations on O‘ahu, Moloka‘i, Lāna‘i, Maui, Kaua‘i, and Hawai‘i. Shrubs to tall trees.
<i>incana</i>	Low to middle elevations on O‘ahu, Moloka‘i, Lāna‘i, Maui, and Hawai‘i. Shrubs to tall trees.
<i>macrophylla</i>	Middle elevations on Hawai‘i. Small to tall trees. Relatively large leaves.
<i>newelli</i>	Low to middle elevations on Hawai‘i, usually along watercourses. Small to large trees.
<i>polymorpha</i>	Middle to high elevations on O‘ahu, Moloka‘i, Lāna‘i, Maui, and Hawai‘i. Small to large trees. Leaves usually with dense woolly pubescence on lower surface.
<i>pumila</i>	Middle to higher elevation swamps on Kaua‘i, Moloka‘i, and Maui. Prostrate shrubs.
<i>pseudorugosa</i>	Higher elevation bogs on West Maui. Prostrate shrubs.

are restricted in geographic distribution across the main Hawaiian islands.

ASSOCIATED PLANT SPECIES

‘Ōhi‘a is by far the dominant forest tree in Hawai‘i and makes up most of the biomass in most lowland and montane wet and mesic forests. ‘Ōhi‘a also occurs as a major component of the vegetation in dry forests, subalpine shrublands, and on new lava flows. ‘Ōhi‘a forests in wet areas with deep soil are tall in stature with a continuous canopy, whereas scattered ‘ōhi‘a trees occur on new or little-weathered lava flows and in dry areas. Small ‘ōhi‘a trees grow with other shrub species at high-elevation sites above 2000 m (6500 ft). Open stands occupy young lava flows in wet areas on Hawai‘i island, while swampy areas may be occupied by dwarf ‘ōhi‘a trees, especially on the island of Kaua‘i (Adee and Conrad 1990, Wagner et al. 1999).

Associated species commonly found in native habitats

‘Ōhi‘a forms almost pure stands on new lava flows. In older forests it grows as huge old specimens along with up to 40 other tree species (Lamb 1981). In wet and mesic areas above 800 m (2600 ft), where most remaining native forests occur, the next most common tree after ‘ōhi‘a is koa (*Acacia koa*). ‘Ōhi‘a and koa share the montane forests with smaller tree species such as naio (*Myoporum sandwicense*), kōlea (*Myrsine lessertiana*), ‘ōlapa (*Cheirodendron* spp.), pilo (*Coprosma* spp.), kāwa‘u (*Ilex anomala*), manono (*Hedyotis* spp.), alani (*Melicope* spp.), kōpiko (*Psychotria* spp.), and ‘ōhi‘a hā (*Syzygium sandwicense*). The understory is dominated by ferns, including the tree ferns hāpu‘u (*Cibotium* spp.) and ‘ama‘u (*Sadleria* spp.), and shrubs such as the native Hawaiian raspberry or ‘ākala (*Rubus harwaiensis*) and ‘ōhelo (*Vaccinium* spp.). Dryland ‘ōhi‘a forests are interspersed with native trees such as lama (*Diospyros sandwicensis*), alahe‘e (*Psydrax odorata*), kauila (*Colubrina oppositifolia*), olopuia (*Nestegis sandwicensis*), wiliwili (*Erythrina sandwicensis*), and naio. Lama, alahe‘e, and hala (*Pandanus tectorius*) are also common in the few remaining lowland wet ‘ōhi‘a forests, where the understory is frequently the uluhe fern (*Dicranopteris linearis*). At high elevations, above about 2000 m (6500 ft), ‘ōhi‘a forms scattered stands and shrublands with māmane (*Sophora chrysophylla*), koa, and naio (Adee and Conrad 1990, Wagner et al. 1999).

Species commonly associated as aboriginal introductions in Pacific islands

Few coastal ‘ōhi‘a forests remain, and ‘ōhi‘a is rarely seen in cultivated areas in Hawai‘i, which were originally cleared for agriculture and now support the Polynesian-introduced trees such as coconut, breadfruit (*Artocarpus altilis*), and milo (*Thespesia populnea*).

Species commonly associated in modern times or as recent introductions

Native ‘ōhi‘a forests in Hawai‘i suffer from an onslaught of invasive alien species. Chief among these in the wet forests are strawberry guava (*Psidium cattleianum*), albizia (*Falcataria moluccana*), and various melastomes (for example, *Melastoma* spp., *Clidemia hirta*, and *Miconia calvescens*).



Heidi Johansen stands next to a centuries-old ‘ōhi‘a lehua tree that has a circumference at breast height of 6.8 m (22.3 ft) and diameter of 2.16 m (7.08 ft). Manukā, South Kona, Hawai‘i. PHOTO: C. ELEVITCH

‘Ōhi‘a forests in mesic areas suffer invasions from *Morella faya* (formerly *Myrica*), *M. cerifera*, and Christmas berry (*Schinus terebinthifolius*). The understory of ‘ōhi‘a forests is often composed of alien grasses, in particular meadow rice grass (*Ehrharta stipoides*), which inhibits regeneration of the trees. Dryland ‘ōhi‘a forests are invaded by silk oak (*Grevillea robusta*) and threatened by fires encouraged by invasion of the introduced fountain grass (*Pennisetum setaceum*).

LEGENDARY 'ŌHI'A LEHUA

Many Hawaiians believe that 'ōhi'a forests are sacred to Pele, the goddess of volcanoes. When angry, Pele destroys the 'ōhi'a with streams of lava. It is one of five plants sacred to Laka, the goddess of hula. Many Hawaiian legends include 'ōhi'a.

In one legend Ka-ehu, the yellow shark of Pearl Harbor, homesick for the beauty of Puna, chanted:

*O my land of rustling lehua-trees!
Rain is treading on your budding flowers,
It carries them to the sea.
They meet the fish in the sea.
This is the day when love meets love,
My longings are stirring within me
For the spirit friends of my land.
They call me back to my home, I must return.*
(Westervelt 1916)

ENVIRONMENTAL PREFERENCES AND TOLERANCES

Climate

'Ōhi'a has a wide environmental amplitude. The species occurs from just above sea level to the tree line (above 2500 m [8200 ft]), where such high-altitude forests are exposed to frost and ephemeral snows. Along moisture gradients 'ōhi'a can be found distributed between sites as extreme as dry forests with less than 400 mm (16 in) annual rainfall to wet forests and bogs with more than 10,000 mm (33 ft) annual rainfall (Stemmermann and Ihsle 1993, Dawson and Stemmermann 1999). Maximum biomass and optimal growth form occur at mean annual temperatures between 16°C (61°F) and 21°C (70°F) and rainfall between 1000 (40 in) and 3000 mm (120 in) annually (Stemmermann 1983; Stemmermann and Ihsle 1993; Mueller-Dombois 1987, 1994; Adee and Conrad 1990).

Elevation range

1–2500 m (3–8200 ft)

Mean annual rainfall

400–10,000 mm (16–400 in); 1000–3000 mm (40–120 in) preferred

Rainfall pattern

Prefers year-round rainfall, although it grows in areas with seasonal precipitation.

Dry season duration (consecutive months with <40 mm [1.6 in] rainfall)

4 months

Mean annual temperature

10–24°C (50–75°F); 16–21°C (61–70°F) preferred

Mean maximum temperature of hottest month

29°C (84°F)

Mean minimum temperature of coldest month

0°C (32°F)

Minimum temperature tolerated

-4°C (25°F)

Soils

The range of soil types tolerated includes Histosols, Mollicsols, Spodosols, Oxisols, Ultisols, and Alfisols (Adee and Conrad 1990, Crews et al. 1995). 'Ōhi'a is the dominant tree on new, excessively drained lava flows as well as on ancient and water-saturated soils surrounding the Alaka'i bogs on Kaua'i (Mueller-Dombois 1994, Herbert and Fownes 1995). 'Ōhi'a is also dominant on both nutrient-rich and nutrient-depleted soils (Crews et al. 1995, Herbert and Fownes 1999), as well as soils suffering from aluminum toxicity problems (Moomaw et al. 1959, Mueller-Dombois 1994). On exposed ridges, steep slopes, or poorly drained sites 'ōhi'a does not reach large size and may be reduced to dwarf shrub stature (Adee and Conrad 1990).

Soil texture

'Ōhi'a usually grows on medium and heavy clay soils of the Hawaiian uplands. It also tolerates rocky soils and organic soils forming on new lava flows.

Soil drainage

The species develops best on relatively level well drained sites. 'Ōhi'a does not tolerate poor drainage well and may die out on very poorly drained pāhoehoe (sheet) lava.

Soil acidity

pH 3.6–7.4. 'Ōhi'a generally grows in acid, weathered soils where the pH may reach as low as 3.6 (Crews et al. 1995). In drier areas 'ōhi'a may grow in soils that are neutral to mildly alkaline.

Special soil tolerances

'Ōhi'a tolerates shallow, rocky soils, and young trees may be found on all but the very freshest lava flows. In other sites, 'ōhi'a will grow in extremely acid, infertile soils.



'Ōhi'a is often one of the first plants to colonize new lava flows. PHOTO: C. ELEVITCH



Dwarf 'Ōhi'a near the Alaka'i swamp, Kaua'i, Hawai'i. In harsh conditions, or on poorly drained sites, 'Ōhi'a is dwarfed. PHOTO: D. HERBERT

Tolerances

Drought

Well established 'Ōhi'a with deep root systems can persist in extremely dry areas receiving less than 400 mm (16 in) of rain annually. On older substrates in dry areas, 'Ōhi'a forests are replaced by other native dryland species such as lama (*Diospyros sandwicensis*) and alahe'e (*Psydrax odorata*) (Stemmermann and Ihsle 1993). Droughts in these areas commonly last for 4 months. However, trees growing on shallow lava soils in wetter areas can be killed by droughts of a few weeks.

Full sun

'Ōhi'a prefers to grow in full sun.

Shade

The tree tolerates only light shade. It can grow underneath a koa canopy but does not regenerate in its own shade or under other understory trees, relying on treefall gaps in the mature forest.

Fire

'Ōhi'a is killed by intense fires, although it may resprout after light fires. Dryland 'Ōhi'a forests are threatened by alien invasive grasses, which allow fires to spread and kill the native trees (Smith and Tunison 1992).

Frost

'Ōhi'a tolerates light frosts, but temperatures below -8°C (18°F) permanently damage tissue (Scowcroft et al. 2000, Cordell et al. 2000).

Waterlogging

It does not tolerate waterlogging. Forest dieback is common in high-rainfall areas with underlying heavy clay soils or pāhoehoe (sheet) lava (Hodges et al. 1986).

Salt spray

The species has low salt tolerance and is seldom found where exposed to salt spray, although it is found in sheltered areas down to sea level (Rauch and Hensley 1997).

Wind

'Ōhi'a has good wind tolerance and is wind-firm in hurricanes (Herbert et al. 1999), although strong winds may snap isolated trees.

Other

The tree is tolerant of volcanic fumes.

Abilities

Regenerate rapidly

'Ōhi'a regenerates prolifically from windblown seeds after disturbances, if the site is not taken over by alien vegetation. 'Ōhi'a is the only native tree to regenerate on recent lava flows for the first few years or even decades.

Self-prune

The tree self-prunes only slowly. Tall trees may retain scraggly lower branches for years.

Coppice

Sometimes 'Ōhi'a coppices thickly, but at other times the tree is killed by coppicing or pollarding. New growth after trees are coppiced is fragile and easily broken. Fallen trees



‘Ōhi‘a trees often start out growing on nurse logs on the forest floor (left)... leaving a cavity under the tree after the substrate decays (right). Pictured: co-author J.B. Friday. PHOTOS: J. B. FRIDAY, T. MCEVOY

in the forest, however, may send up new vertical shoots that in time develop into trees.

GROWTH AND DEVELOPMENT

‘Ōhi‘a trees seed prolifically but grow slowly. They survive by colonizing lava flows or fallen logs in forest gaps where other vegetation cannot compete. Growth of mature forest trees may be imperceptibly slow.

Growth rate

In optimal conditions, seedlings 60–70 cm (24–28 in) tall can be grown from seed in a year. This relatively fast rate of growth may continue through the pole stage, after which growth is typically slow.

Stem diameter growth rates are especially slow, in the range of 1–3 mm (0.04–0.12 in) per year. Saplings and poles may have relatively fast growth rates (Bornhorst 2005), but forest trees rarely grow more than 2 mm (0.08 in) in diameter per year (Adee and Conrad 1990, Herbert and Fownes 1995, 1999, Raich et al. 1997, Vitousek and Farrington 1997, Gerish and Mueller-Dombois 1999). Annual height growth is up to 0.3–0.6 m (12–24 in).

Flowering and fruiting

The earliest flowering is 2 years after germination (Bornhorst 2003). Flowering occurs throughout the year but may be most abundant after seasonal rains, generally during and after the winter months and on into summer. The seed capsules may not release their seed for up to a year after flowering (Carpenter 1976).



Some trees form air roots that trap moisture droplets in the air. PHOTO: C. ELEVITCH

Timber yields

Timber volumes per unit area for mature forest are low in comparison for related trees such as *Eucalyptus* species because of 'ōhi'a's tendency to grow in dense, pole-sized stands and the trees' irregularly shaped stems, which yield little lumber. Higher-volume stands may only yield 70–84 m³/ha (1000–1200 ft³/ac or 5000–6000 bf/ac) (Skolmen 1974a), whereas the average stands yield only half that. Basal area of 'ōhi'a forests may be up to 40 m²/ha (175 ft²/ac), although again this may be composed of many small trees (Adee and Conrad 1990). A forest survey in 1970 estimated that stands on deep soils on the wetter parts of the island, which were described as “commercial” 'ōhi'a stands, were increasing volume at a rate of 0.47 m³/ha/yr (6.7 ft³/ac/yr or 34 bf/ac/yr). Many other stands were composed of pole-sized timbers that were not increasing in volume at all (Metcalf et al. 1970).

Rooting habit

'Ōhi'a is shallow-rooted on pāhoehoe (sheet) lava or in poorly-drained areas but deeply rooted in dry sites and on 'ā'ā (rough) lava. Even on pāhoehoe lava, roots may find cracks and penetrate deeply.

Reaction to competition

'Ōhi'a does not tolerate shading from overstory trees and can easily be smothered or shaded out by exotic weedy grasses, vines, and trees.

PROPAGATION

'Ōhi'a is propagated by cuttings, air-layering, or by seeds. While seeds germinate readily and are true to type (Stemmermann 1983), cuttings or air-layers are perhaps the best way to achieve specific flower color for ornamentals (Rauch and Hensley 1997). The National Tropical Botanical Garden recommends that care be taken to select plants or seeds to be propagated from a nearby locale with similar environmental conditions to the outplanting site, because 'ōhi'a exhibits a variety of growth patterns that are influenced by environment (NTBG 1993).

Propagation by seed

Seeds are easily collected and germinate readily. In natural settings seeds commonly germinate as epiphytes on tree ferns or moss-covered logs (Little and Skolmen 1989).

Seed collection

Whole fruit capsules are collected. Fruits mature 70–90 days after flowering and should be collected after maturity

but before capsules open (Allen 2002). The mature capsules will be swollen, and lines will appear where the capsule will dehisce (Allen 2002). Capsules should be air-dried in a paper bag or cardboard box to prevent loss of seeds (NTBG 1993, Herring 2002). Viability ranges from less than 10% to more than 75% (Dawson 1970, Burton 1982, Drake 1993, Allen 2002).

'Ōhi'a can flower throughout the year but flowering generally peaks in spring or summer after vegetative flushing. Some varieties or populations peak in fall or winter. Individual trees or branches may produce flowers at any time during the year (Adee and Conrad 1990). Drake (1993) noted that on the island of Hawai'i (wet forests at 700 m [2300 ft] elevation), 75% of seeds were dispersed in December and January.



Capsules must be collected before they open, as shown here, or the seeds disappear into the wind. PHOTO: J. B. FRIDAY

Seed storage

For greatest viability, seeds should be sown as quickly as possible after collection (NTBG 1993). Seeds may be stored in a dry, cool location or under refrigeration (Allen 2002), but for greatest viability seeds should be sown within a month of capsule dehiscence. Viability may drop to less than 10% after a year of storage (Allen 2002) and may be near 0% within 3 years (Corn 1979).

Pre-planting treatments

No pretreatment of seeds is necessary.

Germination, media, and containers

Seeds should be spread onto germination trays filled with a moist, sterile potting medium, sterile compost, or cinder and left uncovered or covered by a very thin layer of soil (NTBG 1993, Bornhorst 2005). Fresh seeds germinate in 5–10 days, whereas stored seeds may take as long as 6 weeks

(Corn 1979). Seedlings should be transplanted into 5 cm (2 in) diameter containers with a well drained, loose substrate when 4–6 true leaves have formed (NTBG 1993, Bornhorst and Rauch 2003). The National Tropical Botanical Garden recommended a small amount of slow-release NPK 8–8–8 fertilizer with micronutrients be mixed with the media upon planting, and once-per-month applications of a liquid foliar fertilizer at half strength be applied thereafter (NTBG 1993). Bornhorst (2003) recommended daily watering of potted seedlings.

Time to outplanting

With good nursery practices, seedlings can be ready for outplanting in 6–12 months at a size of 25–30 cm (10–12 in) tall.

Outplanting

Seedlings are vulnerable to drought and should be well established before planting out (NTBG 1993, Bornhorst 2005, Allen 2002). Full sun, frequent watering, and avoidance of strong wind are recommended.

Other comments on propagation

Transplanting of wild seedlings seldom succeeds. If transplanting is attempted, plants should be as small as possible, not more than a few cm tall.

Potted seedlings have a tendency to send roots down into the ground through the bottom of their containers. Care should be taken that they do not form taproots into the ground while in the nursery. This is a particular problem in warm, humid environments.

Propagation by cuttings

Collecting vegetative material

Cuttings can be taken from the tips of vigorous and recently matured growth and should be 1 cm (0.4 in) diameter and 10–15 cm (4–6 in) long (Criley 1998, Bornhorst and Rauch 2003). Success can vary with material selected. Near 100% success is reported from some parent material, while other material is nearly impossible to propagate (Bornhorst and Rauch 2003).

Treatment

Bornhorst (2005) recommended cutting leaves



Top: A container that prevents root spiraling is recommended. Pictured: Baron Horiuchi. Bottom: For certain situations where early field maintenance is impractical, trees can be grown out in large containers and successfully transplanted. Pictured: Anya Tagawa. PHOTOS: C. ELEVITCH

in half, removing leaves from the lower 2.5 cm (1 in) of the stem, and dipping cuttings in a strong rooting hormone for ten seconds. Rauch and Hensley (1997) had 100% success with a 2000 ppm concentration indolbutyric acid (IBA) at a ratio of 2:1 with naphthaleneacetic acid (NAA). Criley (1998) reported success with rooting hormones having 2:1 IBA to NAA at concentrations of 2000–4000 ppm IBA. To grow ‘ōhi‘a from cuttings, a misting system is strongly recommended.

Media

Sterile potting media such as pure perlite or vermiculite, 1:1 perlite to vermiculite, or 2:1 mixture of perlite to peat moss have been used (Bornhorst 2005, Rauch and Hensley 1997, Criley 1998, Herring 2002). Choice depends on climate and grower’s preference.

Propagation by air-layers

Vegetative material

Some individuals may air-layer easier than others (Herring 2002). Bornhorst (2005) suggested that the presence of aerial roots might indicate a plant that will air-layer easily.

Treatment

Herring (2002) recommended using a standard air-layering technique, while Bornhorst (2005) recommended using a strong rooting hormone. Tanabe and Frazier (1984) had

good success with the ring-girdle technique and a range of indolbutyric acid IBA concentrations.

DISADVANTAGES

Potential for invasiveness

Because of its prolific, windblown seeds and ability to survive on harsh sites, ‘ōhi‘a has the potential to become a serious pest if planted outside the Hawaiian islands. On other Pacific islands, there is the danger that ‘ōhi‘a would hybridize with native populations of other species of *Metrosideros*.

Diseases and pests

The National Tropical Botanical Garden (NTBG) notes problems with thrips and ants, which can transfer other pests such as scale and aphids to ‘ōhi‘a (NTBG 1993). These pests mainly affect young, planted trees. Leaves on young trees are frequently affected by psyllid insects that cause galls. Mealybugs and sooty mold may affect seedlings in wet areas. A previously undescribed rust was found on ‘ōhi‘a nursery seedlings in early 2005 (D. Ogata, pers. com.). The endemic cerambycid borer *Plagithmysus bilineatus* has the greatest potential for negative impact on ‘ōhi‘a. It may become epidemic to weakened trees in cases of canopy dieback (Papp et al. 1979, Mueller-Dombois 1985, Adee and Conrad 1990). Other potentially damaging borers include

Ceresium unicolor, *Xyleborus saxesensi*, and *X. simillimus* (Adee and Conrad 1990). ‘Ōhi‘a is attacked by the two spotted leafhopper (*Sophonia rufofascia*), an exotic pest only discovered in Hawai‘i in 1987 (Alyokhin et al. 2004). Forest trees are host to several species of native Hawaiian mistletoe (*Korthalsella* spp.) (Wagner et al. 1999). Root rots *Phytophthora cinnamomi*, *Pythium vexans*, and *Armillaria mellea* can also be damaging and are most often associated with weakened stands experiencing canopy decline (Corn 1972, Papp et al. 1979). ‘Ōhi‘a forests on the windward side of the island of Hawai‘i were observed to be undergoing massive canopy dieback in the 1960s and 70s. Five to seven types of decline have been identified, and no single uniform cause has been determined. Most of the area affected by ‘ōhi‘a decline is wet and poorly drained, and the so-called “wetland dieback” and “bog formation dieback” make up more than 80% of the affected decline area (Hodges et al. 1986). While the fungal rots *Phytophthora cinnamomi* and *Armillaria mellea* and the borer



Air-layering is frequently used to clone a desirable tree, such as this one with yellow blossoms. PHOTO: C. ELEVITCH

A NEW DISEASE OF 'ŌHI'A

A new rust disease was discovered on potted 'ōhi'a plants in a nursery on O'ahu in April 2005. No previous rusts had been reported for 'ōhi'a, thus this discovery was a cause for concern. The rust has been identified as *Puccinia psidii* and causes a common disease of eucalyptus, guava, allspice (*Pimenta dioica*), paperbark (*Melaleuca quinque-nervia*), and other plants in the Myrtaceae in South and Central America to Florida, where it is called eucalyptus rust or guava rust. The rust quickly spread to all the major islands and on O'ahu has been found infecting rose apple (*Syzygium jambos*) and common guava as well as the native trees *Eugenia koolauensis* and *E. reinwardtiana*.

The rust most commonly affects young shoots. Initial symptoms are yellow powdery spots growing individually or in a circular pattern on the leaf and shoot. Infected young tissue does not expand and leaves and shoots become deformed as normal tissue continues to expand. As this is a new disease in Hawai'i, it is unknown how severe the problem will be in natural forests or whether the rust can kill mature 'ōhi'a trees. *Puccinia psidii* is considered a serious disease in the eucalyptus industry in Brazil. The Hawai'i Department of Agriculture is asking that grow-

ers avoid transporting 'ōhi'a, eucalyptus, paperbark, guava, rose apple, or other seedlings in the family Myrtaceae inter-island to reduce the spread of the disease, although the rust spores are wind-borne and naturally spread rapidly within a community. For ornamental trees, cutting and destroying infected foliage and shoots may help reduce the infection. In nurseries, keeping the foliage dry may also help prevent infection (Killgore and Heu 2005).



Rust symptoms on new shoot. PHOTO: D. OGATA

Plagithmysus bilineatus have all been found attacking trees in areas affected by dieback, all probably only affect trees already stressed by poor soil drainage or other environmental factors. The most likely explanation of the 'ōhi'a dieback phenomenon is that 'ōhi'a stands are predisposed to decline because of age and cohort senescence (Mueller-Dombois et al. 1980, Mueller-Dombois 1983). Mature 'ōhi'a stands are even-aged in nature, having been regenerated after a disturbance, and therefore reach senescence at the same time and are then vulnerable to attacks of pests and diseases. Prolific 'ōhi'a regeneration under most dieback stands lends credence to the theory that 'ōhi'a decline is age-related. In some areas, however, bogs are formed after the dieback of the 'ōhi'a forest.

Other disadvantages or design considerations

'Ōhi'a's chief limitation is its slow growth. Timber plantations of 'ōhi'a are unknown and would probably not be financially viable.

AGROFORESTRY/ENVIRONMENTAL PRACTICES

Soil stabilization

'Ōhi'a forests protect the upper reaches of most of Hawai'i's critical watersheds. On new lava flows, 'ōhi'a forests help break down the rock and develop organic soils as well as providing sites for litter accumulation.

Crop shade/overstory

If some 'ōhi'a trees are left when a forest is cleared for agricultural use, these can provide light shade for coffee, cacao, or other shade-loving understory crops. Alternatively, tree crops such as noni (*Morinda citrifolia*) may be planted in alleys cut in the 'ōhi'a forest.

Shade or street tree

'Ōhi'a saplings and poles will grow quickly given an adequate water supply (Bornhorst 2005) but will not develop a significant canopy for many years. Trees are susceptible to dieback if the sites are badly compacted and often flooded and are susceptible to physical damage by people or careless weeding and maintenance.

Homegardens

Because of improved propagation success, 'ōhi'a has in recent years become a popular garden ornamental. The flowers are most often red but are also found in yellow, orange, and salmon. Many garden shops and nurseries now carry 'ōhi'a lehua (Bornhorst 2003, 2005). Unfortunately, some growers sell exotic species of *Metrosideros* as a type of 'ōhi'a (e.g., "New Zealand 'ōhi'a") when they are aliens such as pohutukawa (*M. excelsa*) and *M. kermadecensis*. Alien species in the same genus as 'ōhi'a may pose a threat to the genetic pool of the native Hawaiian species if they escape from cultivation.

Windbreaks

Natural stands of 'ōhi'a make fine windbreaks for moderately windy areas. Single trees or single rows of trees, if left after the surrounding forest is cut, are susceptible to wind breakage, so bands several meters wide should be left. Planted 'ōhi'a trees could be part of a windbreak if planted along with other, faster growing trees.

Silvopasture

Isolated 'ōhi'a trees are often found in high-elevation pastures in Hawai'i, remnants of the forests that were originally cleared. These trees are seldom healthy and suffer from soil compaction and root damage from grazing livestock. Eventually, if an area is grazed long enough, all the 'ōhi'a trees except for individuals growing in protected locations die off. 'Ōhi'a cannot regenerate through thick pasture grasses such as kikuyu grass (*Pennisetum clandestinum*), and cattle damage any young trees that get started.

Watershed and habitat for native species

'Ōhi'a is more valuable for its environmental services than for products. 'Ōhi'a forests are important habitat for Hawai'i's native and endemic species of birds, arthropods, mollusks, and plants, many of which have co-evolved with the 'ōhi'a and endangered (Zimmerman 1948, Hadfield and Mountain 1980, Berger 1981, Hadfield and Miller 1989, Smith et al. 1995). The honeycreeper family of birds, in particular, has evolved into many endemic species in Hawai'i. Most of these, such as the common, bright red 'apapane (*Himatione sanguinea*), the scarlet 'i'iwi (*Vestiaria coccinea*), and the yellow 'amakihi (*Loxops virens*) inhabit the 'ōhi'a forest and feed on the nectar of the 'ōhi'a



'Ōhi'a can survive in pasture but will not regenerate. PHOTO: C. ELEVITCH



Dispersed 'ōhi'a trees can provide light shade for crops such as coffee, as shown here, without interfering with most farm operations. PHOTO: C. ELEVITCH

blossoms and insects in the trees (Carlquist 1980). 'Ōhi'a and koa forests also provide habitat for Hawai'i's most endangered birds, such as the Hawai'i 'ākepa (*Loxops coccinea*), which nests in the cavities of the largest 'ōhi'a trees (Freed 2001), and the Hawai'i creeper (*Oreomystis mana*). The endangered 'akiapōlā'au (*Hemignathus munroi*) has evolved a woodpecker-like lower bill and the ability to drill holes and

suck sap directly from certain individual ‘ōhi‘a trees (Pejchar, pers. com.). ‘Ōhi‘a also provides critical specific habitat for endemic *Achatinella* and *Partulina* tree snail species (Hadfield and Mountain 1980, Hadfield and Miller 1989). Understory plants in the forest germinate on fallen ‘ōhi‘a logs or even on still-living ‘ōhi‘a trees. The trees provide a growing surface above the alien grasses, which cover the forest floor and protect young plants against damage by feral pigs.

‘Ōhi‘a provides valuable watershed protection in Hawai‘i because of its abundance and especially broad environmental amplitude. Most of the upper-elevation forests above Hawai‘i’s important agricultural and urban areas are covered with ‘ōhi‘a forests.

Ornamental

‘Ōhi‘a trees, with their brightly colored flowers and silvery or reddish young foliage, make beautiful ornamental specimen plants.

USES AND PRODUCTS

Honey

‘Ōhi‘a is an excellent honey plant, and ‘ōhi‘a honey is one



The native Hawaiian honeycreeper ‘apapane on an ‘ōhi‘a lehua blossom.
PHOTO: J. JEFFREY

of the only truly native Hawaiian honeys. It is commonly sold in markets throughout Hawai‘i.

Medicinal

In ancient times extracts of ‘ōhi‘a blossoms mixed with other herbs were used to treat childbirth pains and thrush (a fungal disease). In modern times flowers are still given to mothers for childbirth pains (Krauss 2001). New leaf shoots with a reddish color were chewed for numbing sore throats.

Wood

The wood is very hard and dense. A set of tests at the USDA Forest Service Forest Products Laboratory on wood from eleven trees from Kona and Puna found an average specific gravity of 0.81 (oven dry weight over green volume) (Youngs 1960). Weight averaged 0.91 g/cm³ (57 lb/ft³) for oven-dry wood and 1.17 g/cm³ (73 lb/ft³) for green wood. Shrinkage was 6.9% radial, 12.1% tangential, and 19.1% in volume. In color ‘ōhi‘a has pale brown sapwood that merges indistinctly to the reddish to purplish brown heartwood, often streaked, and can develop a long curl. The interlocked grain and growth bands sometimes give the wood an attractive figure; at other times the wood is very plain. The texture is medium and the vessels are very fine and scattered throughout (Lamb 1981). The wood is not as strong as its density would suggest, though, probably because of its spiral grain (Youngs 1960). ‘Ōhi‘a’s hardness



Although ‘ōhi‘a is not considered feasible for commercial plantation forestry, existing ‘ōhi‘a stands can be interplanted with other timber species such as koa. PHOTO: C. ELEVTICH



Left: 'Ōhi'a seedling that volunteered on fence post, showing how opportunistic the tree can be. Right: A mature 'Ōhi'a tree in upper Puna supporting epiphytic growth of pa'īnu (*Astelia menziesiana*) and *Trematolobelia grandifolia*. PHOTOS: A. YEH

makes it difficult to work, and the wood is notorious for shrinking, warping, splitting, and checking during drying. Because of the wood's tendency to warp, boards should be air-dried before kiln-drying (Youngs 1960). The common spiral grain weakens the wood if sawn into timbers. Severe spiral grain and shrinkage during drying leads to instability of the wood and limits 'ōhi'a's use in furniture making, cabinetry, and construction (Skolmen 1974a, Little and Skolmen 1989). Today the most common uses of the wood include wood strip flooring, decking, decorative posts, and round wood construction (Skolmen 1974a, Little and Skolmen 1989). While floors made of 'ōhi'a in Hawai'i are generally stable, shrinking and swelling of the wood might cause problems in highly seasonal climates. Round posts are varnished to preserve the golden brown color of the sapwood; otherwise they weather to grey as does all exposed wood.

Heartwood is not resistant to attack by fungi when in contact with the ground. In tests in a wet area on O'ahu, stakes cut from 'ōhi'a heartwood lasted only about a third as long as redwood stakes and about half as long as stakes

cut from *Eucalyptus robusta*. 'Ōhi'a posts and rails exposed to the weather but not in contact with the soil were resistant to decay over the 9½ year test (Skolmen 1974a, 1974b). 'Ōhi'a is only slightly resistant to termite attack (Grace et al. 1996).

Historically, 'ōhi'a wood has been used as fence posts, furniture, veneer, ukulele keys, ship blocking in dry-docks, bracing stakes in agriculture and irrigation ditch construction, wharf fenders and marine construction, pallets, pile-driver cushions, and biofuel (Skolmen 1974a, Little and Skolmen 1989, Grossman 1992). A mill in Pāhoa on the island of Hawai'i sold millions of railroad ties made of 'ōhi'a to the Santa Fe Railroad company in the early 1900s, until it was found that the ties were not durable in the desert climate.

Ancient Hawaiians used the wood in carving sacred images (ki'i or tikis), spears, and mallets, and household items such as poi boards. Construction uses included structural poles, rafters, and temple walls (Abbott 1992, Gon and Pang 1998, Kirch 1985, Little and Skolmen 1989).

Fence posts

‘Ōhi‘a poles are traditionally used as fence posts in Hawai‘i, although they are only moderately durable when in contact with the soil, and longer lasting woods such as kauila (*Colubrina oppositifolia* or *Alphitonia ponderosa*) were preferred before they became rare.

Fuelwood

‘Ōhi‘a has always been a favorite firewood, continuing to today. It burns with a hot, clean flame. The only disadvantage is the difficulty in splitting the cross-grained logs.

Craft wood

Jewelry, jewelry boxes, turned bowls, furniture, and cabinetry are among the many modern craft uses. ‘Ōhi‘a was not used for carved wooden calabashes (‘umeke lā‘au) in ancient times in Hawai‘i due to its tendency to warp and crack (Abbott 1992).

Canoe/boat/raft making

Despite the large size of ‘ōhi‘a trees, they were not used to build hulls for canoes (wa‘a) in ancient times. Once a canoe’s hull was carved from a koa tree, ‘ōhi‘a wood was often used for the decking, gunwales, and seats (Abbott 1992). Spreaders for canoe hulls were carved from naturally curved ‘ōhi‘a stilt roots (Krauss 1993). ‘Ōhi‘a wood was also sometimes used for paddles and for the booms that connected the hulls of the great double-hulled voyaging canoes (wa‘a kaulua).

Body ornamentation/garlands

The flowers are used in garlands and leis and in the traditional Hawaiian religion are considered sacred to the goddess Pele (Kirch 1985, Little and Skolmen 1989, Bornhorst 2005). The tips of young shoots known as liko are also commonly incorporated into leis.

Ceremonial/religious importance

The majority of the large carved wooden sacred images (ki‘i) in old Hawai‘i were made from the wood of ‘ōhi‘a. Along with luma (*Diospyros sandwicensis*), ‘ie‘ie (*Freyinetia arborea*), halapepe (*Pleomele aurea*), maile (*Alyxia oliviformis*), and the palapalai fern (*Microlepia strigosa*), ‘ōhi‘a was one of the ritual plants associated with the hula. ‘Ōhi‘a represented the god Kūkā‘ōhi‘a Laka, who was named for a legendary ‘ōhi‘a tree that bore both red and



Top: ‘Ōhi‘a wood flooring. PHOTO: J. B. FRIDAY **Bottom: Hawaiians frequently use the wood for sacred images (ki‘i), which turn light gray when exposed to the elements. Pu‘uhonua o Hōnaunau, Kona, Hawai‘i.** PHOTO: C. ELEVITCH

white flowers (Abbott 1992). In the traditional Hawaiian religion, the fire-red flowers are considered sacred to Pele, the volcano goddess.

Other

The red flowers are the official symbol of the island of



Liko (the new leaf tips) are found in a rainbow of colors and are a favorite component of leis. PHOTOS: C. ELEVITCH

Hawai'i. Some people believe that picking 'ōhi'a flowers will cause it to rain.

URBAN AND COMMUNITY FORESTRY

'Ōhi'a can be a splendid ornamental tree in cooler, wetter areas such as the Hilo and Puna districts on Hawai'i or in Mānoa or Wahiawā on O'ahu. Planting an 'ōhi'a tree in a yard or garden brings a bit of the forest home. A good example of this can be seen in Waimea on Hawai'i, where 'ōhi'a was planted along the roads and in many public landscapes. The tree grows slowly and needs to be well cared for. It cannot tolerate abuse as other popular landscape trees can. If forest areas are cleared for development, specimen 'ōhi'a trees can be left as a reminder of the forest that once was there.

Size

'Ōhi'a trees in landscape settings usually reach 3–12 m (10–40 ft) in height, smaller than trees in the forest. Some trees grow vertically and may only have a canopy spread of 1–2 m (3.3–6.6 ft); others may fork and spread their canopy over 10 m (33 ft).

Rate of growth in a landscape

In well watered areas, 'ōhi'a grows about 60 cm (2 ft) in height per year.

Roots

When grown on lava soils, 'ōhi'a has spreading surface roots. The tree may be killed if these roots are cut, and traffic and machinery can also damage roots. In deep soils or in porous lava, roots go deep and enable the tree to survive dry periods.

Products commonly used in a Hawaiian household

The red, yellow, and orange 'ōhi'a flowers are made into intricate leis in Hawai'i. The silvery, green, or reddish new shoots are also used. 'Ōhi'a wood was once used for housing and carving sacred images (ki'i) in old Hawai'i. Today 'ōhi'a wood is mostly used as round posts (unsawn) for architectural elements and for flooring.

Light requirements

'Ōhi'a trees prefer full sunlight but tolerate slight shading from surrounding plants.

Water/soil requirements

'Ōhi'a prefers wetter and cooler areas in Hawai'i. The tree can grow on various textures of soil and on organic soils formed over lava rock. 'Ōhi'a is killed by drying out and must be kept well watered, especially when first planted (Bornhorst 2005). 'Ōhi'a trees planted in shallow depressions in pāhoehoe (sheet) lava may flourish for several years but die from drying out when the trees grow larger and the

thin soil is no longer able to hold enough water to support them during dry spells. 'Ōhi'a growing on pāhoehoe may also die from constant flooding, especially if water is diverted to the site because of nearby construction or development.

Expected life span in a homegarden

Huge old 'ōhi'a trees in the forest are many centuries old, and even moderate-size trees may be a couple of hundred years old. However, trees grown near places where people live often succumb to injury, root damage, flooding, or soil compaction if care is not taken.

Varieties favored for use in a homegardens or for street trees

Many people favor the mamo (yellow-flowered) variety of 'ōhi'a. Pink, orange, and red are also planted in landscaping. All colors may be propagated by air-layering.

Seasonality of leaf flush, flowering, fruiting

'Ōhi'a may flower at any time of year. Individual trees flower once or twice per year, and some individuals may flower almost continuously.

Exceptional ornamental values

'Ōhi'a is a favorite ornamental tree because of its showy blossoms. Trees with blossoms of exceptional color may be

propagated by cutting or air-layer to ensure that the colors remain true.

Use as living fence, hedge, or visual/noise barrier

'Ōhi'a does not tolerate lopping or pollarding well, and this limits its use as a hedge. However, closely planted trees can form a thick barrier in open spaces.

Bird/bee/wildlife

'Ōhi'a trees attract bees and other insects. While they are important nectar trees for native forest birds, few of these survive near inhabited areas, and 'ōhi'a trees are more likely to attract alien bird species.

Maintenance requirements

Trees should be mulched heavily and fertilized only sparingly. 'Ōhi'a trees have evolved to survive in infertile soils; heavy fertilization causes sappy growth that may not be strong enough to bear its own weight and is prone to insect attack. Slow-release fertilizers or natural compost works best.

Drawbacks

In a landscape setting, 'ōhi'a is limited by its slow growth. Trees over 22–30 cm (10–12 in) tall are not easily transplanted, and larger potted specimens may not survive. 'Ōhi'a is also easily barked and damaged by machinery such as mowers and weed cutters. 'Ōhi'a generally does not tolerate topping or pollarding. A drawback to using 'ōhi'a in



Left: 'Ōhi'a trunk and roots are easily de-barked by careless use of grass trimmers or other machinery. PHOTO: J. B. FRIDAY Center and right: Plastic guards can help protect bark from damage during maintenance, although damage can still occur, as here above and below guard. PHOTOS: C. ELEVTICH



Left: Stand of 'ōhi'a serving as a windbreak for the Hawai'i Belt Road on the Hāmākua coast. PHOTOS: J. B. FRIDAY Right: Gall insects commonly affect young trees, but the trees usually outgrow these pests. PHOTO: C. ELEVITCH

agroforestry systems is that it is a host for the black twig borer (*Xylosandrus compactus*), a serious pest of coffee, cacao, and other tree crops.

Nuisance issues

None.

Common pest problems

Young 'ōhi'a foliage may be attacked by Chinese rose beetles and gall insects. Usually the trees outgrow these pests without treatment. Aphids and their associated ants and sooty mold may be problems in wet areas. A rust disease probably caused by the fungus *Puccinia psidii* was discovered in Hawai'i in 2005 (see text box above).

Other

Developers in Hawai'i often leave a few token 'ōhi'a when clearing native forest for building sites. Unfortunately, these usually die after a few months or years from having their root system damaged by bulldozers and other machinery. If 'ōhi'a are desired as part of the landscape, an area at least as wide as the trees are tall must be left undisturbed. Since many building sites on Hawai'i are on relatively new lava flows with shallow soils, the trees' root systems are spreading and shallow. Cutting or trampling the roots will usually kill the trees. Compromised trees can become a hazard, especially in high winds, as the whole tree can come crashing down.

COMMERCIAL PRODUCTS

'Ōhi'a lehua has become a popular garden ornamental, and many nurseries and garden shops now carry the tree

(Bornhorst 2003, 2005). For production of ornamental flowers it is recommended to start plants from cutting or air-layer because vegetative propagation ensures that the plants will be true to type (Rauch and Hensley 1997).

'Ōhi'a posts are sold in Hawai'i today for approximately US\$1.00 per inch (2.5 cm) of diameter per lineal foot (30 cm). For example, a ten-foot long post of eight inches in diameter would sell for \$80.00. Air-dried, rough-sawn lumber suitable for milling into flooring sells for \$1.25/bf, while lumber of firsts and seconds grade sells for \$3.50–\$5.00 per board foot. Finished 'ōhi'a flooring sells for \$7.00–\$9.00 per square foot.

Spacing

Because of its slow growth rate, 'ōhi'a is not grown in timber plantations.

Management techniques

Timber management is seldom if ever undertaken in 'ōhi'a forests because of the tree's perceived low value and slow rate of growth. Faster-growing pole stands might be managed for pole production for architectural elements. At a growth rate of 3 mm (0.12 in) diameter per year, it would take 50 years to grow a 15 cm (6 in) diameter pole. Natural forests might be extensively managed for a small return. However, continued pressure to clear lowland forests for agricultural crops makes the prospect of such sustainable, long-term silviculture unlikely.

Design considerations

While 'ōhi'a itself is slow growing and unlikely to yield much return, 'ōhi'a trees can be managed as part of a koa silvicultural system, as occurs in the native forests. The

faster-growing koa does not shade out the 'ōhi'a, but the 'ōhi'a can provide side shade for the koa and possibly reduce branchiness and encourage straight growth (Patrick Baker, pers. com.). 'Ōhi'a could also provide habitat for native birds in such a mixed system. Koa trees may be planted into an existing scattered 'ōhi'a forest, or 'ōhi'a may be planted along with koa. 'Ōhi'a is not shade tolerant and cannot be regenerated under its own canopy or in dense shade, although it does grow well under a light koa canopy.

On-farm processing methods required to access market

'Ōhi'a is often logged and milled using portable mills brought to the forest sites. Most 'ōhi'a harvesting today occurs in conjunction with land clearing for coffee, papaya, pasture, or other agriculture or residential uses. Almost all 'ōhi'a harvested is cut on the island of Hawai'i.

On-farm processing

'Ōhi'a logs often check badly at the ends. Brackets hammered into the ends of logs may help decrease losses due to splitting. Drying lumber should be done very slowly to minimize checking and splitting (Forest Products Laboratory, undated). Poles should be debarked when freshly cut.

Markets

'Ōhi'a flooring is produced locally on O'ahu and Hawai'i. 'Ōhi'a architectural elements are favored by builders on Hawai'i and upcountry areas on Maui and Kaua'i.

INTERPLANTING/FARM APPLICATIONS

Example 1

Location

Hakalau National Wildlife Refuge, Hawai'i island. The refuge is located on the upper slopes on the windward side of Mauna Kea, in a cool, mesic area. Rainfall is 2000 mm (80 in) annually. The deep, acid soils are derived from volcanic ash and classified as Typic Hydrudands (USDA classification). Soils are silty clay loams and silt loams which contain high amounts of organic material but also fix available phosphorus. Temperatures at the higher elevations can dip below

freezing during the winter months (Scowcroft and Jeffrey 1999). While originally forested, the upper parts of the refuge had been gradually converted by logging, burning, and grazing to pasture dominated by kikuyu grass (*Pennisetum clandestinum*), meadow rice grass (*Ehrharta stipoides*), and other alien grasses over the past century.

Description

The U.S. Department of the Interior's Fish and Wildlife Service has been restoring a native koa-'ōhi'a forest to the higher elevations (above 1800 m [6000 ft]) of the refuge since 1989. Their goal is to restore the habitat for the native bird life, including eight endangered species such as the 'ākepa (*Loxops coccinea*), the 'akiapōlā'au (*Hemignathus munroi*), and the Hawai'i creeper (*Oreomystis mana*). 'Ōhi'a trees are grown from locally-collected seed in a nursery at the site. The seedlings, which are grown in dibble tubes



'Ōhi'a seedlings planted under 8- to 10-year-old koa trees at Hakalau National Wildlife Refuge are protected from frost damage. PHOTO: C. ELEVITCH

or root trainers, take 18–24 months to reach plantable height of 15–20 cm (6–8 in) tall. Planting sites are scalped to remove competing grasses, and trees are initially fertilized with 10–30–10 granular fertilizer. Older trees at the site are growing at a rate of approximately 1 mm (0.05 in) per year in diameter. Early reforestation efforts indicated that a major cause of mortality in young outplanted ‘ōhi‘a seedlings was frost damage. Currently ‘ōhi‘a seedlings are outplanted underneath the canopies of 8–10-year-old koa trees established on the site. The koa overstory decreases radiative cooling at ground level and keeps the ‘ōhi‘a seedlings from being damaged (Scowcroft et al. 2000). Koa, being a nitrogen-fixing tree, also increases nitrogen availability in the soil. When planted under koa, ‘ōhi‘a trees have shown close to 100% survival, whereas they showed almost no survival when planted on bare ground in the open.

An alternative successful planting technique is to plant seedlings onto fallen logs in the pastures, or even onto artificial planters made from wood and potting material. Elevating the seedlings above the ground by even 0.5 m (20 in) seems to help avoid frost damage.

Spacing

The ‘ōhi‘a seedlings are planted at wide spacings, about 4 m (13 ft) apart, in a mixture of other native forest and understory trees including kōlea (*Myrsine lessertiana*), ‘ōlapa (*Cheirodendron* spp.), and pilo (*Coprosma* spp.), and the shrubs ‘ākala (*Rubus hawaiiensis*), pūkiawe (*Leptecophylla tameiameia*), and ‘ōhelo (*Vaccinium* spp.).

Example 2

Location

Puna, Hawai‘i.

Description

‘Ōhi‘a forests on lava rock soils in lower Puna are often cleared for planting horticultural crops such as papaya, noni, or ‘awa (kava). These crops all benefit from windbreaks. While planting ‘ōhi‘a trees is not practical because of their slow growth, some landowners carefully lay out their fields so as to leave remnants of the ‘ōhi‘a forest to serve as windbreaks.

Crop/tree interaction

The crops benefit from decreased moisture stress in dry periods and avoid physical damage from the wind. Windbreaks of ‘ōhi‘a also serve

as honey plants for nearby beekeepers and as forage for any native birds that remain in the area.

Spacing

Windbreaks generally protect areas ten times their height. ‘Ōhi‘a forests in the lava rock lands of Puna may only be 10 m (40 ft) tall. To protect 3 m (10 ft) tall papaya or other tree crops, windbreaks should be left every 100 m (330 ft). Since ‘awa and noni tolerate some shade, windbreaks may be spaced even closer for better protection, or plantings may be done in alleys cleared in the native forest. Since a single row of ‘ōhi‘a may not be windfirm, especially on shallow substrates, enough land needs to be left so that the trees can grow several rows deep.

Example 3 (Contributed by Craig Elevitch)

Location

Kona, Hawai‘i.

Description

Coffee is traditionally grown in full sun in Kona, but it can also be grown under light shade. Some landowners choose to leave some of the native forest trees such as ‘ōhi‘a and lama (*Diospyros sandwicensis*) as shade trees in coffee farms. While both these trees are slow growing and it would not be practical to plant them along with the coffee, it is sometimes possible to leave some trees in place when the farm is cleared. These trees may provide habitat for native birds and serve as a reminder of the history of the farm. One disadvantage is that old ‘ōhi‘a trees, as any nearby trees, may serve as a host for the black twig borer (*Xylosandrus compactus*), a serious pest of coffee.



Windbreak of ‘ōhi‘a forest protecting a noni plantation in Puna, Hawai‘i.

PHOTO: S. NELSON

Crop/tree interactions

Although coffee is a shade-tolerant plant, shading decreases the amount of the coffee produced. Light overstory shade is generally used in tropical countries to decrease overbearing and consequent nutrient stress (Bittenbender and Smith 2004). While growers in Hawai'i generally have avoided nutrient stress by fertilizing sufficiently and growing coffee in full sun, there may be a marketing premium or niche for "shade-grown" coffee.

Spacing

'Ōhi'a and other native overstory trees are left as they grow, scattered across the landscape.

PUBLIC ASSISTANCE

The Cooperative Extension Service (CES) of the University of Hawai'i can assist landowners with questions relating to 'ōhi'a. Their forestry web site includes many valuable publications, forestry news, and an extensive list of forestry links for Hawai'i.

Extension Forester

College of Tropical Agriculture and Human Resources

University of Hawai'i at Mānoa

Komohana Agricultural Complex

875 Komohana St., Hilo, HI 96720

Tel: 808-959-9155; Fax: 808-959-3101

Web: <http://www.ctahr.hawaii.edu/forestry>

INTERNET

Hardwood Drying Schedules, Forest Products Laboratory, USDA Forest Service: <<http://www.fpl.fs.fed.us/drying.html>>.

Hawaiian Native Plant Propagation Database—*Metrosideros polymorpha*: <<http://www2.hawaii.edu/~eherring/hawnprop/met-poly.htm>>.

Native Plant Network Propagation Protocol Database: <<http://www.nativeplantnetwork.org/network/search.asp>>.

The Flora of the Hawaiian Islands, a web site of the Smithsonian Institution, is a database of flowering plants and ferns in Hawai'i: <<http://ravenel.si.edu/botany/pacificislandbiodiversity/hawaiianflora/index.htm>>.



Coffee planted in the understory of native 'ōhi'a-lama-kolea forest, 'Ōhi'a Forest Farm, South Kona, Hawai'i. PHOTO: C. ELEVITCH

BIBLIOGRAPHY

(☛ indicates recommended reading)

Abbott, I.A. 1992. *Lā'au Hawai'i: Traditional Hawaiian Uses of Plants*. Bishop Hawai'i Museum Press, Honolulu.

Adams, N.M. 1967. *New Zealand Native Trees*. Reed Ltd., Wellington, New Zealand.

Adee, K., and C.E. Conrad. 1990. *Metrosideros polymorpha* Gaud. In: R.M. Burns and B.H. Honkala (eds.). *Silvics of North America. Agricultural Handbook 654*. USDA Forest Service, Washington, DC.

- Allen, J.A. 2002. *Metrosideros polymorpha* Gaudich. In: Vozzo, J.A. (ed.). Tropical Tree Seed Manual. Agricultural Handbook 721. United States Department of Agriculture, Washington, DC.
- Alyokhin, A.V., P. Yang, and R.H. Messing. 2004. Oviposition of the invasive two-spotted leafhopper on an endemic tree: Effects of an alien weed, foliar pubescence, and habitat humidity. *Journal of Insect Science* 4: 13. <<http://www.insectscience.org/4.13>>.
- Aradhya, K.M., D. Mueller-Dombois, and T.A. Ranker. 1991. Genetic evidence for recent and incipient speciation in the evolution of Hawaiian *Metrosideros* (Myrtaceae). *Heredity* 67: 129–138.
- Berger, A.J. 1981. Hawaiian Birdlife. 2nd Ed. University of Hawai'i Press, Honolulu.
- Bittenbender, H.C., and V.E. Smith. 2004. Growing Coffee in Hawai'i, rev. ed. College of Tropical Agriculture and Human Resources, University of Hawai'i, Honolulu.
- Bornhorst, H. 2003. Hawai'i Gardens: Native 'ohi'a lehua plant a perfect fit for Hawai'i gardens. Honolulu Advertiser May 23, 2003.
- Bornhorst, H.L., and F. Rauch. 2003. Native Hawaiian Plants for Landscaping, Conservation, and Reforestation. Ornamentals and Flowers Series No. 30. College of Tropical Agriculture and Human Resources, University of Hawai'i, Honolulu.
- Bornhorst, H.L. 2005. Growing Native Hawaiian Plants: A How-to Guide for the Gardener, rev. ed. Bess Press, Honolulu.
- Burton, P.J. 1982. The effect of temperature and light on *Metrosideros polymorpha* seed germination. *Pacific Science* 36: 229–240.
- Carlquist, S. 1980. Hawaii: A Natural History. National Tropical Botanical Garden, Lāwā'i, Hawai'i.
- Carpenter, F.L. 1976. Plant-pollinator interactions in Hawaii: Pollination energetics of *Metrosideros collina* (Myrtaceae). *Ecology* 57 (6): 1125–1144.
- Cordell, S.C., G. Goldstein, P.J. Melcher, and F.C. Meinzer. 2000. Photosynthesis and freezing avoidance in ohia (*Metrosideros polymorpha*) at treeline in Hawaii. *Arctic, Antarctic, and Alpine Research* 32 (4): 381–387.
- Corn, C.A. 1979. Variation in Hawaiian *Metrosideros*. Ph.D. Dissertation, University of Hawai'i, Honolulu.
- Corn, C.A., and W.M. Hiesey. 1973. Altitudinal variation in Hawaiian *Metrosideros*. *American Journal of Botany* 60: 991–1002.
- Crews, T.E., K. Kitayama, J.H. Fownes, R.H. Riley, D.A. Herbert, D. Mueller-Dombois, and P.M. Vitousek. 1995. Changes in soil phosphorus fractions and ecosystem dynamics across a long chronosequence in Hawaii. *Ecology* 76(5): 1407–1424.
- Criley, R.A. 1998. Propagation of indigenous and endemic ornamental Hawaiian plants. *Combined Proceedings of the International Plant Propagators' Society* 48: 669–674.
- Dawson, J.W. 1970. Pacific capsular Myrtaceae. 2. The *Metrosideros* complex: *M. collina* group. *Blumea* 18: 441–445.
- Dawson, J.W., and L. Stemmermann. 1999. *Metrosideros* Banks ex Gaertn. In: Wagner, Herbst, and Sohmer, op. cit.
- Drake, D. 1993. Germination requirements of *Metrosideros polymorpha*, the dominant tree of Hawaiian lava flows. *Biotropica* 25: 461–467.
- Evenhuis, N.L., and L.G. Eldredge. 2004. Records of the Hawaii Biological Survey for 2003. Part II. Notes. Occasional Paper 79. Bishop Museum, Honolulu.
- Forest Products Laboratory. undated. WWW Hardwood Drying Schedules. <<http://www.fpl.fs.fed.us/drying.html>>.
- Freed, L.A. 2001. Significance of old-growth forest to the Hawai'i 'ākepa. *Studies in Avian Biology* 22: 173–184.
- Gerrish, G., and D. Mueller-Dombois. 1999. Measuring stem growth rates for determining age and cohort analysis of a tropical evergreen tree. *Pacific Science* 53 (4): 418–429.
- Gon III, S., and B.K. Pang. 1998. The Forest Travels on the Wind. Moanalua Gardens Foundation Hawaiian Culture internet archive. <<http://www.pixi.com/~mgf/archive.html>>.
- Grace, J.K., D.M. Ewart, and C.H.M. Tome. 1996. Termite resistance of wood species grown in Hawaii. *Forest Products Journal* 46(10): 57–61.
- Grossman, D.H. 1992. Early Recovery of a Hawaiian Lowland Rainforest Following Clearcutting at Kalapana on the Island of Hawaii. Ph.D. Dissertation, University of Hawai'i, Honolulu.
- Hadfield, M.G., and B.S. Mountain. 1980. A field study of a vanishing species: *Achatinella mustelina* (Gastropoda: Pulmonata), in the Waianae mountains of Oahu. *Pacific Science* 34: 345–358.
- Hadfield, M.G., and S.E. Miller. 1989. Demographic study of *Partulina proxima*. *Pacific Science* 43: 1–16.
- Herbert, D.A., and J.H. Fownes. 1995. Phosphorus limitation of forest leaf area and net primary productivity on a highly weathered soil. *Biogeochemistry* 29: 223–235.
- Herbert, D.A., and J.H. Fownes. 1999. Forest productivity and efficiency of resource use across a chronosequence of tropical montane soils. *Ecosystems* 2: 242–254.
- Herbert, D.A., J.H. Fownes, and P.M. Vitousek. 1999. Hurricane damage to a Hawaiian forest: nutrient supply rate affects resistance and resilience. *Ecology* 80: 908–920.
- Herring, E. 2002. Hawaiian Native Plant Propagation Database—*Metrosideros polymorpha*. <<http://www2.hawaii.edu/~eherring/hawnprop/met-poly.htm>>.
- Hodges, C.S., K.T. Adee, J.D. Stein, H.B. Wood, and R.B. Doty. 1986. Decline of Ohia (*Metrosideros polymorpha*)

- in Hawaii: a review. General Technical Report PSW-86. USDA Forest Service Pacific Southwest Forest and Range Experiment Station, Berkeley, California.
- James, S.A., C.F. Puttock, S. Cordell, and R.P. Adams. 2004. Morphological and genetic variation within *Metrosideros polymorpha* (Myrtaceae) populations from different habitats on Hawai'i. *New Zealand Journal of Botany* 42: 263–270.
- Killgore, E.M., and R.A. Heu. 2005. A rust disease on ohia. New Pest Advisory 05-04. Hawaii State Department of Agriculture, Honolulu. <<http://www.hawaiiag.org/hdoa/npa/npa05-04-ohiarust.pdf>>
- Kinsey, M. undated. Pohutukawa—the New Zealand Christmas Tree. Global Garden, Australia. <<http://www.global-garden.com.au/backissuez/0112feature1.htm>>.
- Kirch, P.V. 1985. Feathered Gods and Fishhooks. University of Hawai'i Press, Honolulu.
- Kitayama, K., R. Pattison, S. Cordell, D. Webb, D. Mueller-Dombois. 1997. Ecological and genetic implications of foliar polymorphism in *Metrosideros polymorpha* Gaud. (Myrtaceae) in a habitat matrix on Mauna Loa, Hawaii. *Annals of Botany* 80: 491–497.
- Krauss, B.H. 1993. Plants in Hawaiian Culture. University of Hawai'i Press, Honolulu.
- Krauss, B.H. 2001. Plants in Hawaiian Medicine. Bess Press, Honolulu.
- Lamb, S.H. 1981. Native Trees and Shrubs of the Hawaiian Islands. Sunstone Press, Santa Fe, New Mexico.
- Little, E.L., Jr., and R.G. Skolmen. 1989. Common Forest Trees of Hawaii (Native and Introduced). *Agricultural Handbook* 679. USDA Forest Service, Washington, DC.
- Metcalf, M.E., R.E. Nelson, E.Q.P. Petteys, and J.M. Berger. 1970. Hawaii's Timber Resources. Resource Bulletin PSW-15. USDA Forest Service Pacific Southwest Forest and Range Experiment Station, Berkeley, California.
- Moomaw, J.C., M.T. Nakamura, G.D. Sherman. 1959. Aluminum in some Hawaiian plants. *Pacific Science* 13: 335–341.
- Mueller-Dombois, D. 1983. Canopy dieback and successional processes in Pacific forests. *Pacific Science* 37(4): 317–325.
- Mueller-Dombois, D. 1985. 'Ōhi'a dieback in Hawaii: 1984 synthesis and evaluation. *Pacific Science* 39: 150–170.
- Mueller-Dombois, D. 1987. Forest dynamics in Hawaii. *Trends in Evolution and Ecology* 2: 216–220.
- Mueller-Dombois, D. 1994. Vegetation dynamics and evolution of *Metrosideros polymorpha* in Hawaii. *Phytocoenologia* 24: 609–614.
- Mueller-Dombois, D., and F.R. Fosberg. 1998. *Vegetation of the Tropical Pacific Islands*. Springer-Verlag, New York.
- Mueller-Dombois, D., J.D. Jacobi, R.G. Cooray, and N. Balakrishnan. 1980. Ohia Rainforest Study: Ecological Investigations of the Ohia Dieback Problem in Hawaii. Miscellaneous Publication 183. College of Tropical Agriculture and Human Resources, University of Hawai'i, Honolulu.
- National Tropical Botanical Garden (NTBG). 1993. 'Ōhi'a lehua. In: Native Hawaiian plant information sheets. Hawaii Plant Conservation Center, National Tropical Botanical Garden, Lāwa'i, Hawai'i.
- National Tropical Botanical Garden (NTBG). 1996. Ten native Hawaiian trees for urban landscapes. Education and Plant Science Departments. National Tropical Botanical Garden, Lāwa'i, Hawai'i.
- Papp, R.P., J.T. Kleijunas, R.S. Smith Jr., and R.F. Scharpf. 1979. Associations of *Plagithmysus bilineatus* (Coleoptera: Cerambycidae) and *Phytophthora cinnamomi* with decline of 'ohi'a forests in the Island of Hawaii. *Forest Science* 25: 187–196.
- Raich, J.W., A.E. Russell, and P.M. Vitousek. 1997. Primary productivity and ecosystem development along an elevational gradient on Mauna Loa, Hawai'i. *Ecology* 78: 707–721.
- Rauch, F.D., and D. Hensley. 1997. Ohia lehua. *Ornamentals and Flowers Series no. 11*. College of Tropical Agriculture and Human Resources, University of Hawai'i, Honolulu. <<http://www.ctahr.hawaii.edu/oc/freepubs/pdf/OF-11.pdf>>.
- Rock, J. 1974. *The Indigenous Trees of the Hawaiian Islands*. Charles E. Tuttle Co., Inc., Tokyo.
- Scowcroft, P.G., and J. Jeffrey. 1999. Potential significance of frost, topographic relief, and *Acacia koa* stands to restoration of mesic Hawaiian forests on abandoned rangeland. *Forest Ecology and Management* 114: 447–458.
- Scowcroft, P.G., F.C. Meinzer, G. Goldstein, P.J. Melcher, and J. Jeffrey. 2000. Moderating night radiative cooling reduces frost damage to *Metrosideros polymorpha* seedlings used for forest restoration in Hawaii. *Restoration Ecology* 8(2): 161–169.
- Skolmen, R.G. 1974a. Some Woods of Hawaii: Properties and Uses of 16 Commercial Species. Technical Report PSW 8/1974. USDA Forest Service Pacific Southwest Forest and Range Experiment Station, Berkeley, California.
- Skolmen, R.G. 1974b. Natural Durability of Some Woods Used in Hawaii. Research Note PSW-292. USDA Forest Service Pacific Southwest Forest and Range Experiment Station, Berkeley, California.
- Smith, C.W., and J.T. Tunison. 1992. Fire and alien plants in Hawai'i: Research and management implications for native ecosystems. In: Stone, C.P., C.W. Smith, and J.T. Tunison (eds.). *Alien Plant Invasions in Native Ecosystems of Hawai'i*. University of Hawai'i Cooperative National Park Resources Studies Unit, Honolulu.
- Smith, T.B., L.A. Freed, J.K. Lepson, and J.H. Carothers. 1995. Evolutionary consequences of extinctions in popula-

- tions of a Hawaiian honeycreeper. *Conservation Biology* 9: 107–113.
- St. John, H. 1979. *Metrosideros polymorpha* (Myrtaceae) and its variations. *Hawaiian Plant Studies* 88. *Phytologia* 42: 215–218.
- Stemmermann, L. 1983. Ecological Varieties of 'Ōhi'a Varieties (*Metrosideros polymorpha* Myrtaceae), the Dominants in Successional Communities of Hawaiian Rainforest. Ph.D. Dissertation. University of Hawai'i, Honolulu.
- Stemmermann, L., and T. Ihsle. 1993. Replacement of *Metrosideros polymorpha*, 'ōhi'a, in Hawaiian dry forest succession. *Biotropica* 25: 36–45.
- Tanabe, M.J., and B. Frazier. 1984. The influence of auxin on the aird layering of ohia-lehua. In: Rauch, F.D., W.W. McCall, and K.W. Leonhardt (eds.). Second fertilizer & ornamentals short course. Cooperative Extension Service, College of Tropical Agriculture and Human Resources, University of Hawai'i, Honolulu.
- Vitousek, P.M., and H. Farrington. 1997. Nutrient limitation and soil development: experimental tests of a biogeochemical theory. *Biogeochemistry* 37: 63–75.
- Wagner, W.L., D.R. Herbst, and S.H. Sohmer. 1999. *Manual of the Flowering Plants of Hawai'i*, rev. ed. University of Hawai'i Press and Bishop Museum Press, Honolulu.
- Westervelt, W.D. 1916. *Hawaiian Legends of Ghosts and Ghost-Gods: VIII. A Shark Punished at Waikiki*. Ellis Press, Boston. <<http://www.sacred-texts.com/pac/hlog/hlog00.htm>>.
- Youngs, R.L. 1960. Physical, Mechanical, and Other Properties of Five Hawaiian Woods. Forest Products Laboratory Report 2191. USDA Forest Service, Madison, Wisconsin.
- Zimmerman, E.C. 1948. *Insects of Hawaii*. Vol. 1. Introduction. University of Hawai'i Press, Honolulu.



Species Profiles for Pacific Island Agroforestry (www.traditionaltree.org)

Metrosideros polymorpha ('ōhi'a)

Authors: J. B. Friday¹ and Darrell A. Herbert²

1. Cooperative Extension Service, College of Tropical Agriculture and Human Resources, University of Hawai'i, 875 Komohana Street, Hilo, Hawai'i 96720-2757, USA; <<http://www.ctahr.hawaii.edu/forestry>>.



Cooperative Extension Service
College of Tropical Agriculture and Human Resources
University of Hawai'i at Manoa

2. Boston University College of General Studies, 871 Commonwealth Ave., Boston, MA 02215, USA; Tel: 617-358-0320; Fax: 617-353-5868.

Acknowledgments: The authors and publisher thank Susan Cordell, Dale Evans, Clyde Imada, Heidi Johansen, Mark Kimball, Lyman Perry, Forest and Kim Starr, Janice Uchida, and Aileen Yeh for their input. The authors thank Craig Elevitch for contribution of the example of ohia on a coffee farm. Photo contributions from Craig Elevitch, Tom McEvoy, Jack Jeffrey, Scot Nelson, Desmond Ogata, Forest and Kim Starr, and Aileen Yeh are greatly appreciated.

Recommended citation: Friday, J.B., and D.A. Herbert. 2006. *Metrosideros polymorpha* ('ōhi'a), ver. 3.2. In: Elevitch, C.R. (ed.). Species Profiles for Pacific Island Agroforestry. Permanent Agriculture Resources (PAR), Hōlualoa, Hawai'i. <<http://www.traditionaltree.org>>.

Sponsors: Publication was made possible by generous support of the United States Department of Agriculture Western Region Sustainable Agriculture Research and Education (USDA-WSARE) Program; SPC/GTZ Pacific-German Regional Forestry Project; USDA Natural Resources Conservation Service (USDA NRCS); Kaulunani, an Urban Forestry Program of the DLNR Division of Forestry and Wildlife and the USDA Forest Service; State of Hawai'i Department of Land & Natural Resources Division of Forestry & Wildlife; USDA Forest Service Forest Lands Enhancement Program; and Muriel and Kent Lighter. This material is based upon work supported by the Cooperative State Research, Education, and Extension Service, U.S. Department of Agriculture, and Agricultural Experiment Station, Utah State University, under Cooperative Agreement 2002-47001-01327.

Series editor: Craig R. Elevitch

Publisher: Permanent Agriculture Resources (PAR), PO Box 428, Hōlualoa, Hawai'i 96725, USA; Tel: 808-324-4427; Fax: 808-324-4129; E-mail: par@agroforestry.net; <<http://www.agroforestry.net>>. This institution is an equal opportunity provider.

Reproduction: Copies of this publication can be downloaded from <<http://www.traditionaltree.org>>. This publication may be reproduced for noncommercial educational purposes only, with credit given to the source. © 2006 Permanent Agriculture Resources. All rights reserved.

